

Diagram 3. Showing dynamics of changes in the number of oocysts per g droppings in chicks of Group 1 (Subgroups A, B) and the control group C in Series I. The means of graphic presentation are explained in Chapter dealing with "Material and methods".

Discussion

The only means of effective immunization of poultry against coccidiosis of caeca known so far is the infecting of susceptible individuals with appropriate doses of *E. tenella* oocysts. This method certainly reduces the sustained losses owing to the abortive course of the disease, but it does not remove the source of the infection (distributor of oocysts). From this arises the suggestion that further investigations are required to search for methods of weakening the pathogenic properties of coccidia used as immunizing material.




Diagram 4. Some pathological changes in caeca of experimental chicks in Series I (Groups 1, 2). The means of graphic presentation are explained in Chapter dealing with "Material and methods".

It is known (Horton-Smith 1962, Pastuszko 1973 a, b) that fowls become resistant to all-host specific coccidian species only when oocysts of the respective species have provoked a subclinical form of coccidiosis. No cross-immunity between different coccidian species has been recorded. The immunity obtained by natural means is of the nature of total or partial immunization. So it either completely inhibits the endogenous development of coccidia or only prevents development of certain stages of the life cycle. Thus, the final effect is either complete prevention of infection or its abortive asymptomatic course. Most often in practice the latter course of events has been recorded in poultry farms,

Table III

Mean body-weight gain in chicks of the Series II experiments

| Group of chicks | Mean body-weights (g) recorded on experimental days | | | | | |
|------------------|---|---------|----------|-----------|-----------|-----------|
| | 1st day | 7th day | 14th day | 21st day* | 28th day* | 33rd day* |
| 1 | 37 | 56 | 85 | 104 | 142 | 239 |
| 2 | 35 | 48 | 86 | 113 | 175 | 269 |
| 3 | 37 | 54 | 82 | 117 | 191 | 320 |
| 4 | 37 | 56 | 84 | 117 | 172 | 318 |
| Control | 37 | 54 | 86 | 96 | 138 | 195 |
| Free of coccidia | 37 | 56 | 84 | 144 | 255 | 379 |

* Data obtained from groups 1-4 include mean body-weights of birds of both subgroups together




Diagram 5. Some pathological changes in caeca of experimental chicks in Series I (Groups 1-3, and the control group).

and the immunized birds are free of the parasite.

The immunity to coccidiosis has been sufficiently explored and it has been pointed out to schizontogony. The reasons to suppose that the main source of infection is the lymphoid tissue of the caeca and the production of immunity by the birds do not




Diagram 5. Some pathological changes in experimental chicks in Series II (Groups 1-3, and the control group). The means of graphic presentation are explained in Chapter dealing with "Material and methods".

and the immunized birds are natural reservoirs and distributors of the parasite.

The immunity to coccidia, known for a long time, has not been sufficiently explored so far. The mechanism of its production has not been precisely investigated. The results of studies by many authors seem to point to schizogony forms as responsible for the production of immunity to infection of *Eimeria* sp., *E. tenella* above all. There are many reasons to suppose that schizonts of the first and second generations are the main source of antigen. It is also known (Euzéby et al. 1967) that the lymphoid tissue of the bursa Fabricii plays an indubitable part in the production of immunity to coccidiosis in fowls, because after bursectomy the birds do not become resistant. Also the lymphoid hypertrophy

1 chicks in Series I
in Chapter dealing

o 1973 a, b) that
pecies only when
bclinal form of
idian species has
s is of the nature
tely inhibits the
development of
s either complete
urse. Most often
in poultry farms,

ments

| mental days | |
|-------------|-----------|
| 1 day* | 33rd day* |
| 142 | 239 |
| 175 | 269 |
| 191 | 320 |
| 172 | 318 |
| 138 | 195 |
| 255 | 379 |

groups together




Diagram 6. Some pathological changes in experimental chicks in Series II (Groups 1, 2, and the control group). The means of graphic presentation are explained in Chapter dealing with "Material and methods".

of caeca in chicks after immunization and those suffering from coccidiosis produced by fully infective oocysts, as recorded by the present author, points to the importance of the lymphoid cell elements in this respect. The results obtained by many authors (Horton-Smith et al. 1961, Horton-Smith 1962, Burns and Challey 1959, Rose and Long 1970, Pierce et al. 1962, 1963) reveal humoral character of immunity to coccidia in fowls. The sera of immunized animals kill sporozoites and merozoites of the respective *Eimeria* species in vitro. But attempts to transfer immunity to *E. tenella* passively have, until recently, been unsuccessful. No satisfactory results have been obtained either in attempts to induce active immunity in chicks by the use of dead antigen (solution of *E. tenella* schizonts extract).

The recent studies of the effect of ionizing rays on infective parasitic




Diagram 7. Showing the number of coccidia in chicks of Group 1. The means of graphic presentation are explained in Chapter dealing with "Material and methods".

forms began in 1911 along this line in publications have Schwartz 1922, Burr 1949, Alic 1950. These publications show the effect of ionizing rays on coccidia later and so far no




Diagram 7. Showing dynamics of changes in the number of oocysts per g. droppings in chicks of Groups 2-4 (Subgroups A and B) in Series II. The means of graphic presentation are explained in Chapter dealing with "Material and methods".

forms began in 1916, when Tyzzer and Honeij started to work along this line in their studies of nematodes. Since that time many publications have appeared dealing with that problem, to mention Schwartz 1921, Levin and Evans 1942, Alicata and Burr 1949, Alicata 1951, Gould et al. 1953, Soulsby 1961. These publications mainly refer to parasitic nematodes. The studies on the effect of ionizing rays on the organisms of parasitic protozoa began later and so far not many results of these studies have been published




Diagram 8. Showing dynamics of changes in the number of oocysts per g droppings in chicks of Group 1 (Subgroups A and B) in Series II. The means of graphic presentation are explained in Chapter dealing with "Material and methods".

(Albanese and Smetana 1937, Waxler 1941, Hein 1963, Fitzgerald 1965, Mladenović et al. 1966 and others). All the mentioned authors are unanimous in their opinion that the sporulating oocysts of *Eimeria* sp., particularly in the period preceding the completing of sporulation, are more susceptible than non-sporulating oocysts to the activity of ionizing rays. It has also been confirmed that the growth of the dose of radiation is accompanied by a gradual attenuation of the infectivity of oocysts, with a simultaneous preservation of their immunizing properties with regard to susceptible host organisms. The

results of studies Smetana 1937, Mladenović et al. 1966, and others on *E. tenella* oocysts to irradiated oocysts of chicks have, in the present study respect. The present practical conclusion

1. The general *E. tenella*, to the act it difficult and often from the breeding prevention of caeca given to the possible form of artificial

2. It appears from the X-rays of oocysts administered oocysts methods of immunization

3. The preparation of attenuated *E. tenella* oocysts for further detailed studies mentioned oocysts and oocysts to susceptible

Albanese A. A., Smetana 1937, pathogenicity of *E. tenella*

Alicata J. E. 1951, *Parasit.*, 37, 495-501

Alicata J. E., Burns W. C., 1951, effects of radiation on *E. tenella*, 595-596.

Burns W. C., Chalovich 1951, *tenella*. *Expl. Parasit.*

Euzéby J., Caeiro 1963, la coccidiose caecale

cums, de la bourse de

ré-infectés par *E. tenella*

Euzéby J., Caeiro 1963, sur la coccidiose caecale

contre la coccidiose caecale

et de la bourse de la

vét. Méd. comp. Lyon

Fitzgerald P. R. 1965, of *Eimeria* *bovis*, 2, 1-11

Gould S. E., Van Dyke 1963, china larvae. *Am. J. Hyg.*

Hein H. 1963. Vaccination of chickens. *Proc. 17th Int. Conf. Parasit.*

Hein H. 1968. The pathogenicity of *E. tenella* oocysts to susceptible host organisms. *Parasit.*, 22, 1-11.

Horton-Smith C. 1963, the effect of radiation on the infectivity of *E. tenella* oocysts. *Parasit.*, 57, 1-11.

results of studies by certain authors, particularly Albanese and Smetana 1937, Hein 1963 and recently Paškin 1965 and Mladenović et al. 1966, point out that in the case of exposure of *Eimeria tenella* oocysts to X-rays, the dose of radiation should exceed 10,000 R. However, attempts to find the optimum dose of X-rays or the dose of irradiated oocysts which would be the best for practical immunization of chicks have, until recently, been unsuccessful. Thus, the results of the present study seem to provide new valuable information in this respect. The present promising findings suggest the following, mainly practical conclusions.

1. The generally known resistance of coccidia, including *Eimeria tenella*, to the activity of various physical and chemical factors makes it difficult and often quite impossible to eliminate their infective forms from the breeding environment. In the search for methods of effective prevention of caecal coccidiosis in fowls, particular attention should be given to the possibility of obtaining an effective immunizing means in the form of artificially attenuated oocysts.

2. It appears from the results of the present study that the attenuation by X-rays of oocysts exposed to a dose of at least 10,000 R, the dose administered oocysts being sufficiently high, is one of the satisfactory methods of immunization.

3. The preparation of a detailed method of producing suitably attenuated *E. tenella* oocysts for the purpose mentioned earlier, requires further detailed studies on the effect of various doses of X-rays on the mentioned oocysts and on the methods of administering the attenuated oocysts to susceptible chicks.

REFERENCES

- Albanese A. A., Smetana H. 1937. Studies on the effects of X-rays on the pathogenicity of *Eimeria tenella*. *Am. J. Hyg.*, 26, 27-30.
- Alicata J. E. 1951. Effects of roentgen radiation on *Trichinella spiralis*. *J. Parasit.*, 37, 495-501.
- Alicata J. E., Burr G. O. 1949. Preliminary observations on the biological effects of radiation on the life cycle of *Trichinella spiralis*. *Science, N. Y.*, 109, 595-596.
- Burns W. C., Challey J. R. 1959. Resistance of birds to challenge with *Eimeria tenella*. *Expl. Parasit.*, 8, 515.
- Euzéby J., Caeiro V., Pastuszko J. 1967 a. Etudes physio-pathologiques sur la coccidiose caecale du Poulet. 2^e Note: Modifications histologiques des caecums, de la bourse de Fabricius et de la rate de Poussins primo-infectés et ré-infectés par *E. tenella*. *Bull. Soc. Sci. vét. Méd. comp. Lyon*, 69, 197.
- Euzéby J., Caeiro V., Pastuszko J. 1967 b. Etudes physio-pathologiques sur la coccidiose caecale du Poulet. 3^e Note: Essais d'immunisation passive contre la coccidiose caecale par administration de broyats de caecums, de rate et de la bourse de Fabricius prélevés chez des sujets immuns. *Bull. Soc. Sci. vét. Méd. comp. Lyon*, 69, 201-205.
- Fitzgerald P. R. 1968. Effects of ionizing radiation from cobalt-60 on oocysts of *Eimeria bovis*. *J. Parasit.*, 54, 233-240.
- Gould S. E., Van Dyke J. G., Gombert H. J. 1953. Effect of X-rays on trichina larvae. *Am. J. Path.*, 29, 323-337.
- Hein H. 1963. Vaccination against infection with *Eimeria tenella* in broiler chickens. *Proc. 17. Internat. Vet. Congr. (Hannover, Aug. 14-21)*, 2, 1443-1452.
- Hein H. 1968. The pathogenic effects of *Eimeria acervulina* in young chicks. *Expl. Parasit.*, 22, 1-11.
- Horton-Smith C. 1962. Resistencia hacia drogas anticoccidioides inducida

- powiedniego czasu, k
poddane immunizacji
W każdym przypadku
kurczęta czterech gr
oocyst, napromieniono
następnie jedna z po
zywnymi oocystami w
takt z chorymi kurcz
sty o pełnej inwazyj
Kurczęta obu gr
i ciężki, typowy prze
86,4% (Seria I, 40 dn
zowane, z wyjątkiem
dawką 15000 R, wy
niekiedy poronny pr
przez nie oocyst w
z grupami kontrolny
zahamowanie schodz
oocyst w odchodach
zahamowanie endoge
znaczniejsze, im wyż
mieni zastosowana d
łych i zabitych dos
wych w procesie pow
ka zauważyła też, że
w pełni inwazyjnymi
nizowanych niż u kon

W poszukiwaniu metody skutecznego uodparniania kurcząt przeciwko kocydiozie autorka wykonała dwie serie doświadczeń, w których zaraziła do wola dwutygodniowe kurczęta, oocystami *Eimeria tenella*, podanymi uprzednio działaniu różnych dawek promieni X. Kurczęta trzech grup (po 16) serii I otrzymały jednorazowo lub dwukrotnie oocysty, napromieniowane odpowiednio dawkami 10, 20 i 30 tys. R. Po upływie od-

powiedniego czasu kurczęta te, jak również ptaki grupy kontrolnej, nie poddane immunizacji, otrzymały w pełni inwazyjne oocysty *E. tenella*. W każdym przypadku jedna dawka oocyst wynosiła 100 000. W serii II kurczęta czterech grup (po 18) otrzymały jednokrotnie dawkę 75 000 oocyst, napromieniowanych odpowiednio dawkami 15, 20, 25 i 35 tys. R, następnie jedna z podgrup każdej grupy została zarażona w pełni inwazyjnymi oocystami w indywidualnej dawce 140 000, druga zaś przez kontakt z chorymi kurczętami. Grupa kontrolna otrzymała wyłącznie oocysty o pełnej inwazyjności.

Kurczęta obu grup kontrolnych wykazywały spodziewany ostry i ciężki, typowy przebieg kokcydiozy, przy czym śmiertelność wynosiła 86,4% (Seria I, 40 dni obserwacji) i 80% (seria II, 33 dni). Ptaki immunizowane, z wyjątkiem tych, które otrzymały oocysty, napromieniowane dawką 15 000 R, wykazywały w pozostałych grupach obu serii lekkie, niekiedy poronny przebieg kokcydiozy, przy czym liczba wydalanych przez nie oocyst w odchodach była znacznie mniejsza w porównaniu z grupami kontrolnymi. Dwukrotna immunizacja w serii I powodowała zahamowanie schizogonii i gametogonii, co przejawiało się w braku oocyst w odchodach. Ponadto objawy chorobowe były tym słabsze, zaś zahamowanie endogenego cyklu rozwojowego kokcydiów było tym znaczniejsze, im wyższa była, w warunkach doświadczenia, dawka promieni zastosowana do osłabienia oocyst. Badania sekcyjne kurcząt padłych i zabitych doświadczałnie potwierdziły rolę elementów komórkowych w procesie powstawania odporności przeciwko kokcydiozie. Autorka zauważyła też, że średnie przyrosty wagi ciała u kurcząt, zarażonych w pełni inwazyjnymi oocystami były wyraźnie wyższe u ptaków immunizowanych niż u kontrolnych.

zają przeciw-
których za-
tenella, pod-
częta trzech
oocysty, na-
upływie od-